Supplementary Table 3. Baseline characteristics of patients included vs excluded for missing 3-month mRS

Characteristic	Included (n=10,499)	Excluded (n=1,517)	SMD
Age (yr)	70.3±13.2	70.4±13.8	−0.006
Male	6,251 (59.5)	911 (60.1)	−0.010
NIHSS	3 (1–7)	3 (1–8)	−0.095
Hypertension	6,681 (63.6)	952 (62.8)	0.018
Diabetes	3,517 (33.5)	500 (33.0)	0.011
Hyperlipidemia	3,996 (38.1)	632 (41.7)	−0.074
Atrial fibrillation	2,290 (21.8)	326 (21.5)	0.008
Prior antiplatelet	2,581 (24.6)	350 (23.1)	0.035
Prior anticoagulant	1,008 (9.6)	143 (9.4)	0.006
CMB present	5,245 (50.0)	733 (48.3)	0.033
Chronic hemorrhage present	2,474 (23.6)	331 (21.8)	0.042
cSS present	534 (5.1)	71 (4.7)	0.019

Values are presented as mean±standard deviation, number (%), or median (interquartile range). An SMD below 0.10 is conventionally regarded as a negligible between-group difference.
mRS, modified Rankin Scale; SMD, standardized mean difference; NIHSS, National Institutes of Health Stroke Scale; CMB, cerebral microbleed; cSS, cortical superficial siderosis.